

Conformal equivalent metrics in a plane domain are uniquely determined from travel times

Victor .Palamodov

1 Travel time and hodograph

Let E be an Euclidean space with the line element ds and $\Omega \subset E$ be a closed bounded domain with C^1 -boundary Γ . Let $\mathbf{n}$ be a positive C^1 -function in the closure $\bar{\Omega}$ (refraction coefficient); consider the conformal metric $\mathbf{g} = \mathbf{n}^2 ds^2$ in Ω . Take some points $x, y \in \Omega$ and consider a geodesic curve γ between these points. The length $\tau(\gamma)$ of γ is called travel time. It is a multi-valued function $\tau(x, y)$ of the end points x and y of γ , if, at least one geodesic has a pair of conjugated points. This is typical situation, if $\mathbf{n}$ is not constant, moreover, conjugated points appear for any $\mathbf{n} \neq \text{const}$, if $\mathbf{n}$ is constant at infinity, see [5]. The restriction T of the travel time function to $\Gamma \times \Gamma$ is called the *hodograph*. The branches of the hodograph $T(x, y)$ can be distinguished by values of the tangent vector θ at one end, say x . This vector can be found from the differential of the corresponding branch of the hodograph, see Sec.7.

Problem: whether the refraction coefficient $\mathbf{n}$ in Ω is uniquely determined from knowledge of its hodograph T .

In other words, if two conformal metrics in Ω have equal hodographs, does it imply that the metrics coincide?

This problem was studied by Muhometov [10], Muhometov and Romanov [12], Beylkin [3], Bernstein and Gerver [1][2]. The stability of reconstruction of the metric from hodograph was proved in these papers under assumption that the travel time τ is uniquely defined for any pair of points $x, y \in \bar{\Omega}$, that is the hodograph is a usual function. Then the equation $\tau_1(x, y) = \tau_2(x, y)$ for all pairs of points on the boundary implies $\mathbf{n}_1 = \mathbf{n}_2$ in Ω . See also the

papers of Croke [5], Sylvester and Uhlmann [14]. More references are given in the surveys [7] and [13].

We prove here an estimate similar to that of [10] ($n = 2$) without assumption of uniqueness of the travel time.

2 Conformal metric

Take the problem in a more general setting. Let Ω be a closed bounded domain in $E = \mathbb{R}^2$ with the boundary Γ of class C^2 . Let $g = g_{ij}dx^i dx^j$ be a Riemannian metric in Ω of class C^2 ; x^1, x^2 are linear coordinates in E . Let $\mathbf{n} \in C^2(\Omega)$ be a positive function, called refraction coefficient; consider the conformal Riemannian metric $\mathbf{g} = \mathbf{n}g$.

Assumptions on a refraction coefficient: (i) $\mathbf{n} \in C^2(\Omega)$,

(ii) for any point $x \in \Omega$ any geodesics γ that starts at a point $x \in \Omega$ reaches the boundary Γ with a non tangent direction. This implies that there is no waveguide in Ω (non-trapping geometry).

Let $T(E)$ be the tangent bundle on E , $S(\Omega)$ be the bundle of unit circles in $T(\Omega)$ with respect to g . Let $\gamma = \gamma(x, \theta)$ be the geodesic curve that starts from a point $y = y(x, \theta) \in \Gamma$ and arrives at the point x with the unit tangent vector θ . The parameterization $\gamma = \gamma(x, \theta)$, $(x, \theta) \in S(\Omega)$ can be applied to all geodesics in Ω and the function

$$\tau(x, \theta) = \int_{\gamma(x, \theta)} \mathbf{n} ds \quad (1)$$

is defined and is C^1 -continuous on $S(\Omega)$. It vanishes for outgoing geodesics, that is for $\langle \nu | \theta \rangle \leq 0$, where $\nu(x)$ means the inward conormal to Γ at x . For incoming geodesic we have $\langle \nu | \theta \rangle > 0$. So the data of travel times $\tau(x, \theta)$ for $x \in \Gamma$ is the hodograph of the metric $\mathbf{g}$.

3 Differential of the travel time

Let $\mathbf{n}$ be a refraction coefficient in Ω that fulfil the above assumptions (i) and (ii). Fix an arbitrary point $(x, \theta) \in S(\Omega)$; let $\gamma(x, \theta)$ be the geodesic of the metric $\mathbf{g} = \mathbf{n}^2 g$ that arrive at x with the g -unit tangent vector θ and $y \in \Gamma$ be its initial point. Fix a vector θ and consider $\tau(x, \theta)$ as a function of x only.

Proposition 1 *For any point $(x, \theta) \in S(\Omega)$ we have*

$$d_x \tau(x, \theta) = \mathbf{n}(x) \frac{\mathbf{e}(x, \theta)}{\langle \mathbf{e}(x, \theta) | \theta \rangle}. \quad (2)$$

where $\mathbf{e}(x, \theta)$ is a unit covector such that $\langle \mathbf{e}(x, \theta) | \theta \rangle > 0$.

PROOF. Write $d_x \tau(x, \theta) = \lambda \mathbf{n}(x) \mathbf{e}(x, \theta)$ for a unit covector $\mathbf{e}$ and a scalar $\lambda > 0$. We have $\langle d_x \tau(x, \theta) | \theta \rangle = \mathbf{n}(x) > 0$ according to the (1), which implies $\lambda = \langle \mathbf{e}(x, \theta) | \theta \rangle^{-1}$. ►

We have $0 < \langle \mathbf{e}(x, \theta) | \theta \rangle \leq 1$ and can write $\langle \mathbf{e}(x, \theta) | \theta \rangle = \cos \omega$ for some angle $\omega \in (-\pi/2, \pi/2)$.

Choose an orientation in E ; it induces an orientation in the tangent bundle $T(E)$. Take an orthonormal basis (a, b) in the cotangent bundle $T^*(\Omega)$ that is consistent with this orientation. For a tangent vector θ we denote by $\hat{\theta}$ the dual covector; its coordinate expression is $\hat{\theta}_i = g_{ij} \theta^j$. Write this covector in the form $\hat{\theta} = \cos \phi a + \sin \phi b$ for some angle $\phi \in S^1$, which is a coordinate in the bundle of all unit covectors in Ω . The coordinate system x^1, x^2, ϕ is globally defined in this bundle. Let d be the exterior differential in the complex of differential forms on $S^*(\Omega)$; write $d = d_x + d_\phi$, where $d_\phi f = f' d\phi$, $f' \doteq \partial f / \partial \phi$. Differentiating (2) yields

$$dd_x \tau = d_\phi d_x \tau = \mathbf{n}(x) \left[\frac{d_\phi \mathbf{e}(x, \theta)}{\cos \omega} + \frac{\sin \omega}{\cos^2 \omega} d_\phi \omega \wedge \mathbf{e}(x, \theta) \right] \quad (3)$$

Set $\hat{\eta} = -\sin \phi a + \cos \phi b$. The covectors $\hat{\theta}, \hat{\eta}$ form a positively oriented orthogonal frame and satisfy the equations

$$\begin{aligned} \hat{\theta} \wedge \hat{\eta} &= dV, \quad d\hat{\theta} = \hat{\eta} \wedge d\phi, \\ \hat{\eta} \wedge d\hat{\eta} &= \hat{\theta} \wedge d\hat{\theta} = d\phi dV, \end{aligned}$$

where $dV = (\det \{g_{ij}\})^{1/2} dx^1 dx^2$ is the Riemannian volume form in (Ω, g) . Fix the angle ω by the equation $\mathbf{e} = \cos \omega \hat{\theta} + \sin \omega \hat{\eta}$.

4 Comparing two hodographs

Now we estimate difference between two positive functions $\mathbf{n}_1, \mathbf{n}_2$ (refraction coefficients) in terms of hodographs of two conformal metrics $\mathbf{g}_1 =$

$\mathbf{n}_1^2 g, \mathbf{g}_2 = \mathbf{n}_2^2 g$. Denote by $\gamma_1(x, \theta), \gamma_2(x, \theta)$ the corresponding geodesics, by $\tau_1(x, \theta), \tau_2(x, \theta)$ the corresponding travel times. By Proposition 1 we have $d_x \tau_j(x, \theta) = \mathbf{n}_j(x) \cos^{-1} \omega_j \mathbf{e}_j(x, \theta)$ for some angles ω_1, ω_2 such that $\mathbf{e}_j = \cos \omega_j \hat{\theta} + \sin \omega_j \hat{\eta}$. Set $\rho(x, \phi) = \tau_2(x, \theta) - \tau_1(x, \theta)$ and calculate the product

$$\mathbf{R} \doteq d\rho \wedge d_\phi d\rho = (d\tau_2 - d\tau_1) \wedge (d_\phi d\tau_2 - d_\phi d\tau_1)$$

Lemma 2 *We have*

$$\begin{aligned} \mathbf{R} = & \left[\left(\frac{\mathbf{n}_2}{\cos \omega_2} \right)^2 + \left(\frac{\mathbf{n}_1}{\cos \omega_1} \right)^2 - \frac{2 \cos(\omega_1 - \omega_2) \mathbf{n}_1 \mathbf{n}_2}{\cos \omega_1 \cos \omega_2} \right] \hat{\theta} \wedge d\hat{\theta} \\ & + d_\phi (\mathbf{n}_2 \tan \omega_2 - \mathbf{n}_1 \tan \omega_1) (\mathbf{n}_2 - \mathbf{n}_1) \wedge \hat{\theta} \wedge \hat{\eta} \end{aligned} \quad (4)$$

PROOF. Substituting (3) yields

$$\begin{aligned} \mathbf{R} = & \mathbf{n}_2^2 \frac{\mathbf{e}_2 \wedge d\mathbf{e}_2}{\cos^2 \omega_2} + \mathbf{n}_1^2 \frac{\mathbf{e}_1 \wedge d\mathbf{e}_1}{\cos^2 \omega_1} - \mathbf{n}_1 \mathbf{n}_2 \frac{\mathbf{e}_2 \wedge d\mathbf{e}_1 + \mathbf{e}_1 \wedge d\mathbf{e}_2}{\cos \omega_1 \cos \omega_2} \\ & - \mathbf{n}_1 \mathbf{n}_2 \left[\frac{\sin \omega_1 d\omega_1}{\cos^2 \omega_1 \cos \omega_2} + \frac{\sin \omega_2 d\omega_2}{\cos \omega_1 \cos^2 \omega_2} \right] \wedge \mathbf{e}_1 \wedge \mathbf{e}_2. \end{aligned}$$

We have for $\mathbf{e} = \mathbf{e}_j, j = 1, 2$

$$\begin{aligned} \mathbf{e} \wedge d\mathbf{e} &= (\cos \omega \hat{\theta} + \sin \omega \hat{\eta}) \wedge d(\cos \omega \hat{\theta} + \sin \omega \hat{\eta}) \\ &= (\cos \omega \hat{\theta} + \sin \omega \hat{\eta}) \wedge (\cos \omega d\hat{\theta} + \sin \omega d\hat{\eta} - \sin \omega d\omega \hat{\theta} + \cos \omega d\omega \hat{\eta}) \\ &= \cos^2 \omega \hat{\theta} \wedge d\hat{\theta} + \sin^2 \omega \hat{\eta} \wedge d\hat{\eta} + \hat{\theta} \wedge \hat{\eta} d\omega \\ &= \hat{\theta} \wedge d\hat{\theta} + \hat{\theta} \wedge \hat{\eta} d\omega = (1 + \omega') \hat{\theta} \wedge d\hat{\theta}, \end{aligned}$$

since $\hat{\theta} \wedge d\hat{\eta} = d\hat{\theta} \wedge \hat{\eta} = 0$, and

$$\begin{aligned} \mathbf{e}_1 \wedge d\mathbf{e}_2 &= (\cos \omega_1 \hat{\theta} + \sin \omega_1 \hat{\eta}) \wedge d(\cos \omega_2 \hat{\theta} + \sin \omega_2 \hat{\eta}) \\ &= \cos(\omega_1 - \omega_2) \hat{\theta} \wedge d\hat{\theta} + \cos(\omega_1 - \omega_2) \hat{\theta} \wedge \hat{\eta} d\omega_2 \\ &= \cos(\omega_1 - \omega_2) (1 + \omega'_2) \hat{\theta} \wedge d\hat{\theta}, \\ \mathbf{e}_1 \wedge \mathbf{e}_2 &= -\sin(\omega_1 - \omega_2) dV. \end{aligned}$$

Therefore

$$\begin{aligned} \frac{\mathbf{R}}{\hat{\theta} \wedge d\hat{\theta}} &= \mathbf{n}_2^2 \frac{1 + \omega'_2}{\cos^2 \omega_2} + \mathbf{n}_1^2 \frac{1 + \omega'_1}{\cos^2 \omega_1} - \mathbf{n}_1 \mathbf{n}_2 \frac{\cos(\omega_1 - \omega_2)(2 + \omega'_2 + \omega'_1)}{\cos \omega_1 \cos \omega_2} \\ &\quad - \mathbf{n}_1 \mathbf{n}_2 \sin(\omega_2 - \omega_1) \frac{\cos \omega_2 \sin \omega_1 \omega'_1 - \cos \omega_1 \sin \omega_2 \omega'_2}{\cos^2 \omega_2 \cos^2 \omega_1}. \end{aligned}$$

Two terms containing ϕ -derivatives gives

$$\begin{aligned} &\frac{\cos(\omega_1 - \omega_2)(\omega'_2 + \omega'_1)}{\cos \omega_1 \cos \omega_2} + \sin(\omega_2 - \omega_1) \frac{\cos \omega_2 \sin \omega_1 \omega'_1 - \cos \omega_1 \sin \omega_2 \omega'_2}{\cos^2 \omega_2 \cos^2 \omega_1} \\ &= \frac{\omega'_1}{\cos^2 \omega_1} + \frac{\omega'_2}{\cos^2 \omega_2}, \end{aligned}$$

times $\mathbf{n}_1 \mathbf{n}_2$, which finally yields

$$\begin{aligned} \frac{\mathbf{R}}{\hat{\theta} \wedge d\hat{\theta}} &= \left(\frac{\mathbf{n}_2}{\cos \omega_2} \right)^2 + \left(\frac{\mathbf{n}_1}{\cos \omega_1} \right)^2 - \frac{2 \cos(\omega_1 - \omega_2) \mathbf{n}_1 \mathbf{n}_2}{\cos \omega_1 \cos \omega_2} \\ &\quad + \frac{\mathbf{n}_2^2 \omega'_2}{\cos^2 \omega_2} + \frac{\mathbf{n}_1^2 \omega'_1}{\cos^2 \omega_1} - \mathbf{n}_1 \mathbf{n}_2 \left[\frac{\omega'_1}{\cos^2 \omega_1} + \frac{\omega'_2}{\cos^2 \omega_2} \right], \end{aligned}$$

This completes the proof. ►

5 The basic inequality

Theorem 3 *If $\mathbf{n}_1, \mathbf{n}_2$ fulfil the conditions (i) and (ii) and $\mathbf{n}_1 = \mathbf{n}_2$ on Γ , the inequality holds*

$$\int \left(\frac{\mathbf{n}_2}{\cos \omega_2} - \frac{\mathbf{n}_1}{\cos \omega_1} \right)^2 d\phi dV \leq - \int_{S^1} \int_{\Gamma} d_x \rho \wedge d_\phi \rho. \quad (5)$$

PROOF. The last term of (4) is equal to the exact form

$$d \left[(\mathbf{n}_2 \tan \omega_2 - \mathbf{n}_1 \tan \omega_1) (\mathbf{n}_2 - \mathbf{n}_1) \hat{\theta} \wedge \hat{\eta} \right],$$

since $d(\hat{\theta} \wedge \hat{\eta}) = d(a \wedge b) = 0$. It is continuous up the boundary Γ as well as its primitive function $(\mathbf{n}_2 \tan \omega_2 - \mathbf{n}_1 \tan \omega_1) (\mathbf{n}_2 - \mathbf{n}_1)$ in spite of the

functions $\tan \omega_1, \tan \omega_2$ tend to infinity at Γ . This follows from vanishing of $\mathbf{n}_1 - \mathbf{n}_2$ on Γ . Therefore integration over $S(\Omega)$ gives

$$\begin{aligned}
& \int_{S(\Omega)} d\rho \wedge d_\phi d\rho \\
&= \int_{S(\Gamma)} \left[\left(\frac{\mathbf{n}_2}{\cos \omega_2} \right)^2 + \left(\frac{\mathbf{n}_1}{\cos \omega_1} \right)^2 - \frac{2 \cos(\omega_1 - \omega_2) \mathbf{n}_1 \mathbf{n}_2}{\cos \omega_1 \cos \omega_2} \right] d\phi dV \\
&\geq \int_{S(\Gamma)} \left[\left(\frac{\mathbf{n}_2}{\cos \omega_2} \right)^2 + \left(\frac{\mathbf{n}_1}{\cos \omega_1} \right)^2 - \frac{2 \mathbf{n}_1 \mathbf{n}_2}{\cos \omega_1 \cos \omega_2} \right] d\phi dV \\
&= \int_{S(\Gamma)} \left(\frac{\mathbf{n}_2}{\cos \omega_2} - \frac{\mathbf{n}_1}{\cos \omega_1} \right)^2 d\phi dV
\end{aligned}$$

By Stokes'

$$\int_{S(\Omega)} d\rho \wedge d_\phi d\rho = - \int_{S(\Omega)} d(d_x \rho \wedge d_\phi \rho) = - \int_{S(\Gamma)} d_x \rho \wedge d_\phi \rho,$$

since $d_\phi d = -dd_x$, and (5) follows. ►

Corollary 4 *If the hodographs of $\mathbf{n}_1$ and $\mathbf{n}_2$ coincide for all incoming directions (x, θ) , $x \in \Gamma$, then $\mathbf{n}_2 = \mathbf{n}_1$.*

PROOF OF COROLLARY. It is easy to see that the condition implies equation $\mathbf{n}_1 = \mathbf{n}_2$ on Γ . Therefore we can apply Theorem 3 and conclude that $\rho(x, \phi) \equiv 0$. This yields $\mathbf{n}_2 \cos \omega_1 \equiv \mathbf{n}_1 \cos \omega_2$ for all $x \in \Omega$ and unit vectors θ . Fix x and choose θ in such a way that the geodesic $\gamma_1(x, \theta)$ has maximal length. The geodesic γ_1 is orthogonal to Γ at the initial point y_1 and by the Gauss Lemma (see [4]) the end points z of geodesics $\gamma_1(z, \theta)$ such that $\tau_1(z, \theta) = \tau_1(x, \theta)$ run over a curve I that is orthogonal to $\gamma_1(x, \theta)$. Therefore $\mathbf{e}_1 = \hat{\theta}$ and $\omega_1 = 0$. The equation $\cos \omega_1 = 1$ implies $\mathbf{n}_2(x) \leq \mathbf{n}_1(x)$. The opposite inequality is also true. ►

6 Application to the geodesic transform

Corollary 5 *For an arbitrary metric $\mathbf{n}$ that fulfils (i) and (ii) and any real C^1 -smooth function f in Ω that vanishes at Γ we have*

$$\int \frac{d\phi}{\cos^2 \omega} f^2(x) dV \leq - \int_{S^1} \int_{\Gamma} d_x g \wedge d_\phi g \quad (6)$$

where

$$g(x, \theta) = \int_{\gamma(x, \theta)} f ds.$$

PROOF. We may assume that $f \in C^2(\Omega)$ and apply theorem 3 to $\mathbf{n}_1 = \mathbf{n}$, $\mathbf{n}_2 = \mathbf{n} + \varepsilon f$, where ε is a small parameter. The geodesic curves for $\mathbf{n}_1$ and $\mathbf{n}_2$ are the same up to $O(\varepsilon^2)$. Collecting the terms of order ε in (5), we obtain (6). ►

Example. Consider the Euclidean metric $\mathbf{g} = ds^2$ in the unit disc Ω . By a direct calculation

$$\int \frac{d\phi}{\cos^2 \omega} = 2\pi \left(\frac{1+|x|}{1-|x|} \right)^{1/2}, |x| < 1.$$

Write the line integrals g in the standard parameterization $G(p, \varphi) = g(x, \theta)$ of line integrals, where

$$x = (\cos s, \sin s), -\pi/2 \leq s \leq \pi/2, \theta = (\cos \phi, \sin \phi), s - \pi/2 < \phi < s + \pi/2, \\ p = \cos(\psi - \phi), \varphi = \phi + \pi/2$$

and

$$g'_s = -\sqrt{1-p^2}G'_p, \quad g'_\phi = \sqrt{1-p^2}G'_p + G'_\varphi$$

The inequality (6) takes the form

$$\int \left(\frac{1+|x|}{1-|x|} \right)^{1/2} f^2(x) dx \leq - \int_0^{2\pi} ds \int_{s-\pi/2}^{s+\pi/2} g'_s(x, \theta) g'_\phi(x, \theta) d\phi \\ = \int_0^{2\pi} \int_{-1}^1 \left(\sqrt{1-p^2} (G'_p)^2 + G'_p G'_\varphi \right) dp d\varphi.$$

7 Calculation of angular derivative of the travel time

Take an arbitrary refraction coefficient $\mathbf{n}$ in Ω that satisfies (i) and (ii) and determine the angular parameter θ of a ray γ in terms of the hodograph function. Let s be the arc length along Γ that increases clockwise.

Proposition 6 *Let $(x, \theta), x \in \Gamma$ be an arbitrary outward direction and $\gamma(s)$ be the family of geodesic that joins some point $y \in \Gamma$ and $x(s) \in \Gamma$ with outgoing tangent vector $\theta(s)$ defined for $s \in [0, \varepsilon]$ (or for $s \in [-\varepsilon, 0]$) such that $x(0) = x, \theta(0) = \theta$. Then we have*

$$\sin \psi = \frac{1}{\mathbf{n}(x)} \frac{\partial \tau(\gamma(s))}{\partial s} \Big|_{s=0},$$

where ψ is the angle between the outward conormal $-\nu(x)$ and the outward covector $\hat{\theta}$.

PROOF. By the Gauss Lemma the tangent vector θ is orthogonal to the circle of points with the $\mathbf{g}$ -distance $\tau(x, y)$ from y . ►

This formula also holds for any point x conjugate to y in $\gamma(0)$. In particular, $\theta = -\nu$, if $\partial \tau / \partial s = 0$, that is the geodesic $\gamma(0)$ arrives to x with the normal direction $-\nu(x)$. The angle ϕ of θ of the ray $\gamma(x, \theta)$ is equal to $\phi = \psi + \beta + \pi$, where $\beta = \arg \nu$.

References

- [1] I. N. Bernstein, M. L. Gerver, *On the problem of integral geometry for a family of geodesics and on inverse kinematic problem*, DAN SSSR **243** (1978) N2, 303-305 (Russian); Dokl. Akad. Sci. USSR Earth Sci. Sect. **243** (1978).
- [2] I. N. Bernstein, M. L. Gerver, *A condition of distinguishability metrics by travel-time curves*, Methods and algorithms of interpretation of seismological data (Computational seismology **13**), Moscow, Nauka, 1980, pp. 50-73 (Russian).
- [3] G. Ja. Beylkin, *Stability and uniqueness of the solution of the inverse kinematic problem in the multidimensional case*, J. Soviet Math. **21** (1983), 251-254.
- [4] R. L. Bishop, R. J. Crittenden, *Geometry of manifolds*, AMS Chelsea Publishing, 2001.
- [5] C. Croke, *Rigidity and the distance between boundary points*, J. Differential Geom. **33** (1991), N2, 445-464.

- [6] M. L. Gerver, V. M. Markushevich, Dokl. Akad. Nauk SSSR **175** (1967), 334; Dokl. Akad. Nauk SSSR, Earth Science Sec. **175** (1968) 4.
- [7] M. L. Gerver, *New results in the classical travel-time inversion problem*, Appl. Anal. **71** (1999), N1-4, 253-276.
- [8] V. G. Herglotz, *Über der Elastizität der Erde bei Berücksichtigung ihrer variablen Dichte*, J. für Math. Phys. **52** (1905), N3, 257-299.
- [9] R. Michel, *Sur la rigidité imposée par la longueur des géodésiques*, Invent. Math. **65** (1981), 71-83.
- [10] R. G. Muhometov, *Inverse kinematic problem on plane*, Mathematical problems of geophysics, Novosibirsk, VC SO Akad. Nauk SSSR, 1975, **6**, N2, 243-254 (Russian).
- [11] R. G. Muhometov, *The problem of recovery of a two-dimensional Riemannian metric and integral geometry*, Dokl. Akad. Nauk **232** (1977), N1, 27-31.
- [12] R. G. Muhometov, V. G. Romanov, *On the problem of determining an isotropic Riemannian metric in n -dimensional space*, Dokl. Akad. Nauk **243** (1978) N1, 41-44 (Russian); Dokl. Acad. Sci. USSR **243** (1978), 1330-1333.
- [13] P. Stefanov, G. Uhlmann, *Recent progress on the boundary rigidity problem*, Electronic Research Announcements of the American Mathematical Society **11** (2005), 64-70.
- [14] J. Sylvester, G. Uhlmann, *Inverse problems in anisotropic media*, Contemp. Math. **122** (1991), 105-197.